

THE NOTION OF CATEGORY OVER AN ALGEBRAIC STACK

DENNIS GAITSGORY

ABSTRACT. The goal of this note is to spell out the (apparently well-known and intuitively clear) notion of an abelian category over a stack. In the future we will discuss the (much less evident) notion, when instead of an abelian category one considers a triangulated one.

1. Let $\mathcal{C}$ be a $\mathbb{C}$ -linear abelian category. We will assume that $\mathcal{C}$ is closed under inductive limits,¹ i.e., that the tautological embedding $\mathcal{C} \rightarrow \text{Ind}(\mathcal{C})$ admits a right adjoint $\text{limInd} : \text{Ind}(\mathcal{C}) \rightarrow \mathcal{C}$, and that the latter functor is exact. In particular, it makes sense to tensor objects of $\mathcal{C}$ by vector spaces.

2. The affine case. Let A be a commutative algebra. We say that $\mathcal{C}$ is A -linear if we are given a map $A \rightarrow Z(\mathcal{C})$, i.e., if A acts functorially on every $\text{Hom}(X, Y)$ for $X, Y \in \mathcal{C}$. We shall also say that in this case $\mathcal{C}$ "lives over $S = \text{Spec}(A)$ ".

We claim that we have a well-defined functor of tensor product

$$M, X \mapsto M \otimes_A X : A\text{-mod} \times \mathcal{C} \rightarrow \mathcal{C} :$$

If $M = A^I$ for some index set I , then $M \otimes_A X := X^I$, and if $M = \text{coker}(A^I \rightarrow A^J)$, then

$$M \otimes_A X := \text{coker}(X^I \rightarrow X^J),$$

where the (i, j) -entry of the matrix $X^I \rightarrow X^J$ is given by the action of the (i, j) -entry of the matrix $A^I \rightarrow A^J$.

Lemma 3. *The above definition is independent of the presentation of M as a quotient.*

By construction, the functor of tensor product commutes with inductive limits with respect to both M and X , and is right-exact. In addition, we have:

Lemma 4.

(a) *If M is flat, then the functor $X \mapsto M \otimes_A X$ is exact.*

(b) *If $0 \rightarrow M_1 \rightarrow M_2 \rightarrow M_3 \rightarrow 0$ is a short exact sequence of A -modules with M_3 flat, then the sequence*

$$0 \rightarrow M_1 \otimes_A X \rightarrow M_2 \otimes_A X \rightarrow M_3 \otimes_A X \rightarrow 0$$

is also short exact.

(c) *If M is projective and finitely generated, and $M^\vee$ is the dual module, then the above functor admits left and right adjoints, both given by $X \mapsto M^\vee \otimes_A X$.*

¹In what follows by an inductive limit we will mean a limit taken over a small filtering category

Proof. First, let us note that if M is a projective module given by an idempotent of A^I for some set I , then $M \otimes_A X$ is given by the corresponding idempotent of M^I . This implies that the functor of tensor product with a projective A -module is exact. This implies point (a), since every flat A -module can be represented as an inductive limit of projective ones.

Similarly, for point (b) we can assume that M_3 is projective, in which case the short exact sequence splits and the assertion is obvious.

Point (c) is immediate, since we have the adjunctions maps

$$X \rightarrow M^\vee \otimes_A (M \otimes_A X) \simeq M \otimes_A (M^\vee \otimes_A X) \text{ and } M^\vee \otimes_A (M \otimes_A X) \simeq M \otimes_A (M^\vee \otimes_A X) \rightarrow X$$

that satisfy the necessary conditions. $\square$

Finally, we have:

Proposition 5. *Assume that A' is a faithfully-flat algebra over A . Then $A' \otimes_A X \neq 0$ if $X \neq 0$.*

Proof. (Drinfeld)

Lemma 6. *If A' is a faithfully flat algebra over A , then the quotient A'/A is A -flat.*

Clearly, the lemma implies Proposition 5, by Lemma 4(b). $\square$

Proof. (of the Lemma)

It is enough to show that $A'/A \otimes_A A'$ is A' -flat. But

$$A'/A \otimes_A A' \simeq \operatorname{coker}(A' \rightarrow A' \otimes_A A'),$$

and the latter is a split injection. $\square$

We shall say that $X \in \mathcal{C}$ is flat over A (or S) if the functor $M \mapsto M \otimes_A X : A\text{-mod} \rightarrow \mathcal{C}$ is exact.

7. Change of rings. Let $f : \operatorname{Spec}(A') = S' \rightarrow S = \operatorname{Spec}(A)$ is a morphism of affine schemes, corresponding to a homomorphism of algebras $A \rightarrow A'$. There exists a universal A' -linear category $\mathcal{C}'$, which admits an A -linear functor $\mathcal{C} \rightarrow \mathcal{C}'$. We will denote this category by $\mathcal{C} \times_S S'$, and it is constructed as follows:

Objects of $\mathcal{C}'$ are objects $X \in \mathcal{C}$, endowed with an additional action of A' , such that the two actions of A (one coming from $A \rightarrow A'$, and another from $A \rightarrow \operatorname{End}(\mathcal{M})$), coincide. Morphisms in $\mathcal{C}'$ are arrows $X_1 \rightarrow X_2$ in $\mathcal{C}$ that commute with the A' -action.

The functor $\mathcal{C} \rightarrow \mathcal{C}'$ is given by $X \mapsto A' \otimes_A X$, and it will be denoted by f^* . This functor is the left adjoint to the forgetful functor $f_* : \mathcal{C}' \rightarrow \mathcal{C}$.

Set $S'' = S' \times_S S'$, and let $\mathcal{C}''$ denote the corresponding base-changed category over S'' . One naturally defines the category of descent data on $\mathcal{C}'$ with respect to S'' . We will denote it by $\operatorname{Desc}_{S''}(\mathcal{C}')$, and we have a natural functor $\mathcal{C} \rightarrow \operatorname{Desc}_{S''}(\mathcal{C}')$.

Proposition 8. *Suppose that S' is faithfully-flat over S . Then $\mathcal{C} \rightarrow \operatorname{Desc}_{S''}(\mathcal{C}')$ is an equivalence.*

Proof. This is proved by the usual argument, using Lemma 5. $\square$

9. Stacks: approach I. Let $\mathcal{Y}$ be a stack (algebraic in the faithfully flat sense), for which the diagonal morphism $\mathcal{Y} \rightarrow \mathcal{Y} \times \mathcal{Y}$ is affine. This is equivalent to demanding that any morphism $S \rightarrow \mathcal{Y}$, with S an affine scheme, is affine. We are going to introduce the notion of sheaf of abelian categories over $\mathcal{Y}$. In particular, we will obtain a notion of category over a separated scheme.

Let $\text{Sch}_{\mathcal{Y}}^{\text{aff}}$ be the category of affine schemes over $\mathcal{Y}$, endowed with the faithfully flat topology. A sheaf of categories $\mathcal{C}^{sh}$ over $\mathcal{Y}$ is the following data:

- For each $S = \text{Spec}(A) \in \text{Sch}_{\mathcal{Y}}^{\text{aff}}$, a category $\mathcal{C}_S$ over S .
- For $f : S_2 \rightarrow S_1 \in \text{Sch}_{\mathcal{Y}}^{\text{aff}}$, an S_1 -linear functor $f^* : \mathcal{C}_{S_1} \rightarrow \mathcal{C}_{S_2}$, which induces an equivalence $\mathcal{C}_{S_1} \times_{S_1} S_2 \rightarrow \mathcal{C}_{S_2}$.
- For two morphisms $S_3 \xrightarrow{g} S_2 \xrightarrow{f} S_1 \in \text{Sch}_{\mathcal{Y}}^{\text{aff}}$ an isomorphism of functors $g^* \circ f^* \simeq (f \circ g)^*$, such that the natural compatibility axiom for 3-fold compositions holds.

Given a sheaf of categories $\mathcal{C}^{sh}$ over $\mathcal{Y}$ one can form a single category, denoted $\Gamma(\mathcal{Y}, \mathcal{C}^{sh})$ or $\mathcal{C}_{\mathcal{Y}}$ (or simply $\mathcal{C}$, where no confusion is likely to occur) as follows:

Let $S \rightarrow \mathcal{Y}$ be a faithfully flat cover. We define the category $\mathcal{C}_{\mathcal{Y}}$ to be the category of descent data of $\mathcal{C}_S$ with respect to the two maps $S \times S \rightrightarrows S$. Proposition 8 insures that $\mathcal{C}_{\mathcal{Y}}$ is well-defined, i.e., is canonically independent of the choice of the cover S .

Again, by Proposition 8, we have the natural functor $X \mapsto X_S : \mathcal{C}_{\mathcal{Y}} \rightarrow \mathcal{C}_S$ for any $S \in \text{Sch}_{\mathcal{Y}}^{\text{aff}}$, and for $f : S_2 \rightarrow S_1$ a functorial isomorphism $f^*(X_{S_1}) \simeq X_{S_2}$.

When $\mathcal{Y}$ is itself an affine scheme $S = \text{Spec}(A)$, a data of a sheaf of categories $\mathcal{C}^{sh}$ over S is equivalent to a single category over S , which is reconstructed as $\mathcal{C}_S$. In this case we will often abuse the notation and not distinguish between $\mathcal{C}^{sh}$ and $\mathcal{C}_S$.

We will now define a functor

$$\mathcal{F}, X \mapsto \mathcal{F} * X : \text{QCoh}_{\mathcal{Y}} \times \mathcal{C}_{\mathcal{Y}} \rightarrow \mathcal{C}_{\mathcal{Y}}.$$

Let $\mathcal{F}$ be a quasi-coherent sheaf of $\mathcal{Y}$; for $S = \text{Spec}(A) \in \text{Sch}_{\mathcal{Y}}^{\text{aff}}$ we will denote by $\mathcal{F}_S$ the corresponding quasi-coherent sheaf of S . For $X \in \mathcal{C}_{\mathcal{Y}}$ we define

$$(\mathcal{F} * X)_S := \mathcal{F}_S \otimes_A X_S,$$

which by descent gives rise to an object of $\mathcal{C}_{\mathcal{Y}}$.

The above functor has the following properties:

- (i) $\text{QCoh}_{\mathcal{Y}} \times \mathcal{C}_{\mathcal{Y}} \rightarrow \mathcal{C}_{\mathcal{Y}}$ is right exact and commutes with inductive limits.
- (ii) We have a functorial isomorphism $\mathcal{O}_{\mathcal{Y}} * X \simeq X$.
- (iii) We have functorial isomorphisms $\mathcal{F}_1 * (\mathcal{F}_2 * X) \simeq (\mathcal{F}_1 \otimes_{\mathcal{O}_{\mathcal{Y}}} \mathcal{F}_2) * X$, compatible with triple tensor products and the isomorphism of (ii).

By construction, the assertions of Lemma 4 hold in the present context, when we replace $M \otimes_A X$ by $\mathcal{F} \mapsto \mathcal{F} * X$.

10. Descent of categories. Let $f : \mathcal{Y}' \rightarrow \mathcal{Y}$ be a map of stacks, and $\mathcal{C}^{sh}$ a sheaf of categories over $\mathcal{Y}$. It is clear that it gives rise to a sheaf of categories $\mathcal{C}'^{sh} := \mathcal{C}^{sh} \times_{\mathcal{Y}} \mathcal{Y}'$ over $\mathcal{Y}'$, such that for $S \in \text{Sch}_{\mathcal{Y}'}^{\text{aff}}$ the category $\mathcal{C}'_S$ is by definition $\mathcal{C}_S$, where S is regarded as an object of $\text{Sch}_{\mathcal{Y}}^{\text{aff}}$.

If $g : \mathcal{Y}'' \rightarrow \mathcal{Y}'$, it is clear that we have an equivalence of sheaves of categories

$$\mathcal{C}^{sh} \times_{\mathcal{Y}} \mathcal{Y}'' \simeq (\mathcal{C}^{sh} \times_{\mathcal{Y}} \mathcal{Y}') \times_{\mathcal{Y}'} \mathcal{Y}''.$$

Suppose now $\mathcal{C}'^{sh}$ is a sheaf of categories over $\mathcal{Y}'$. Let p_j^i be the projection on the j -th factor from the i -fold Cartesian product $\mathcal{Y}^{(i)}$ of $\mathcal{Y}'$ over $\mathcal{Y}$ to $\mathcal{Y}'$. Let $\mathcal{C}_j^{(i)sh}$ denote the corresponding base-changed sheaf of categories over $\mathcal{Y}^{(i)}$.

Suppose we are given an equivalence of sheaves of categories over $\mathcal{Y}^{(2)} : \mathcal{C}_1^{(2)sh} \simeq \mathcal{C}_2^{(2)sh}$; a natural transformation between the two functors $\mathcal{C}_1^{(3)sh} \rightarrow \mathcal{C}_3^{(3)sh}$, such that the two natural transformations between the two functors $\mathcal{C}_1^{(4)sh} \rightarrow \mathcal{C}_4^{(4)sh}$ coincide.

Proposition 11. *Suppose that $\mathcal{Y}'$ is faithfully flat over $\mathcal{Y}$. Then there exists a well-defined sheaf of categories $\mathcal{C}^{sh}$ over $\mathcal{Y}$ with an equivalence $\mathcal{C}'^{sh} \simeq \mathcal{C}^{sh} \times_{\mathcal{Y}} \mathcal{Y}'$, and which gives rise to the above functors and natural transformations.*

Proof. The assertion readily reduces to the case when both $\mathcal{Y}$ and $\mathcal{Y}'$ are affine schemes, $\text{Spec}(A)$ and $\text{Spec}(A')$, respectively. Let Φ denote the functor $\mathcal{C}_1^{(2)} \rightarrow \mathcal{C}_2^{(2)}$, and T the natural transformation between the functors $\Phi^{1,3}$ and $\Phi^{2,3} \circ \Phi^{1,2}$ between $\mathcal{C}_1^{(3)}$ and $\mathcal{C}_3^{(3)}$.

We define $\mathcal{C}$ to have as objects $X' \in \mathcal{C}'$ endowed with an isomorphism

$$\alpha_{X'} : \Phi((p_2^1)^*(X')) \rightarrow (p_2^2)^*(X'),$$

such that the diagram

$$\begin{array}{ccc} \Phi^{1,3}((p_3^1)^*(X')) & \xrightarrow{T} & \Phi^{2,3} \circ \Phi^{1,2}((p_3^1)^*(X')) \\ \downarrow & & \downarrow \\ (p_3^3)^*(X') & \xleftarrow{\quad} & \Phi^{2,3}((p_3^2)^*(X')) \end{array}$$

commutes. Morphisms in this category are $\mathcal{C}'$ -morphisms, commuting with the data of $\alpha_{X'}$. Evidently, this is an A -linear category.

By construction, we have a functor $\mathcal{C} \rightarrow \mathcal{C}'$, which gives rise to a functor

$$\mathcal{C} \times_{\text{Spec}(A)} \text{Spec}(A') \rightarrow \mathcal{C}'.$$

The fact that the latter is an equivalence is shown by the base-change technique as in the context of quasi-coherent sheaves. $\square$

12. Example: categories with a group-action. Let us consider an example of the above situation, when $\mathcal{Y}' = \text{pt}$, $\mathcal{Y} = \text{pt}/G$, where G is an affine algebraic group. Let $\mathcal{C}'^{sh}$ be a sheaf of categories over $\mathcal{Y}'$, i.e., a plain category. Then the data of an equivalence $\mathcal{C}_1^{(2)} \rightarrow \mathcal{C}_2^{(2)}$ together with a natural transformation as above is what can be reasonably called an action of the group G on $\mathcal{C}'$.

Let us spell this notion out in more detail. We claim that an action of G on a category $\mathcal{C}'$ is equivalent to a data of a functor

$$\text{act}^* : \mathcal{C}' \rightarrow \mathcal{O}_G\text{-mod} \otimes \mathcal{C}',$$

(here $A\text{-mod} \otimes \mathcal{C}'$ denotes the same thing as $\mathcal{C}' \times_{\text{pt}} \text{Spec}(A)$), and two functorial isomorphisms related to this functor. This first isomorphism is between the identity functor on $\mathcal{C}'$ and the

composition $\mathcal{C}' \xrightarrow{\text{act}^*} \mathcal{O}_G\text{-mod} \otimes \mathcal{C}' \rightarrow \mathcal{C}'$, where the second arrow corresponds to the restriction to $1 \in G$.

To formulate the second isomorphism, note that from the existing data we obtain a natural functor

$$\text{act}_A^* : A\text{-mod} \otimes \mathcal{C}' \rightarrow \mathcal{O}_G\text{-mod} \otimes A\text{-mod} \otimes \mathcal{C}' \simeq (\mathcal{O}_G \otimes A)\text{-mod} \otimes \mathcal{C}'$$

for any algebra A .

The second isomorphism is between the two functors $\mathcal{C} \rightarrow \mathcal{O}_{G \times G}\text{-mod} \otimes \mathcal{C}$ that correspond to the two circuits of the diagram

$$\begin{array}{ccc} \mathcal{C}' & \xrightarrow{\text{act}^*} & \mathcal{O}_G\text{-mod} \otimes \mathcal{C}' \\ \text{act}^* \downarrow & & \text{act}_{\mathcal{O}_G}^* \downarrow \\ \mathcal{O}_G\text{-mod} \otimes \mathcal{C}' & \xrightarrow{\text{mult}^*} & \mathcal{O}_{G \times G}\text{-mod} \otimes \mathcal{C}', \end{array}$$

where mult denoted the multiplication map $G \times G \rightarrow G$. These functors must satisfy the usual compatibility conditions.

From Proposition 11, it follows that an action of G on a category $\mathcal{C}'$ is equivalent to the data of a sheaf of categories $\mathcal{C}^{sh}$ over pt/G . (As we shall see later, the latter can be also reformulated as a category with an action of the tensor category $\text{Rep}(G)$.)

By definition, $\mathcal{C} := \mathcal{C}_{\text{pt}/G}$ can be reconstructed as the category of G -equivariant objects of $\mathcal{C}'$. By definition, the latter consists of $X' \in \mathcal{C}'$, endowed with an isomorphism $\alpha_X : \text{act}^*(X') \simeq \mathcal{O}_G \otimes X'$, which is compatible with unit and associativity constraints. Morphisms in the category are $\mathcal{C}'$ -morphisms, compatible with the data of α .

13. Example: categories acted on by a groupoid. Generalizing the above set-up, let S be a base scheme, and $\mathcal{G} \xrightarrow[p_1]{p_2} S$ be an affine groupoid, such that the maps p_1, p_2 (or, equivalently, one of them) are flat. Let $\mathcal{C}^{sh}$ be a sheaf of categories over the quotient stack $\mathcal{Y} = S/\mathcal{G}$. This data can be rewritten as a sheaf of categories $\mathcal{C}'^{sh}$ over S , acted on by $\mathcal{G}$, which means the following:²

We must be given a functor $\text{act}^* : \mathcal{C}' \rightarrow \mathcal{C}' \times_{S, p_1} \mathcal{G}$, which is $\mathcal{O}_S$ -linear if we regard $\mathcal{C}' \times_{S, p_1} \mathcal{G}$ as a category over S via $\mathcal{G} \xrightarrow{p_2} S$, and two functorial isomorphisms related to it. The first isomorphism is a unit constraint, i.e., an isomorphism between the functor

$$\mathcal{C}' \xrightarrow{\text{act}^*} \mathcal{C}' \times_{S, p_1} \mathcal{G} \xrightarrow{1_{\mathcal{G}}^*} \left(\mathcal{C}' \times_{S, p_1} \mathcal{G} \right) \times_{\mathcal{G}, 1_{\mathcal{G}}} S \simeq \mathcal{C}',$$

where $1_{\mathcal{G}} : S \rightarrow \mathcal{G}$ is the unit map.

Formulate the second isomorphism note that for any scheme S' , mapping to S , we obtain a functor

$$\text{act}^* \times \text{id}_{S'} : \mathcal{C}' \times_S S' \rightarrow \mathcal{C}' \times_{S, p_1} (\mathcal{G} \times_{p_2, S} S').$$

²To simplify the notation, we will assume here that S is affine as well.

The second isomorphism is an associativity constraint, i.e., an isomorphism between the two functors in the diagram

$$\begin{array}{ccc} \mathcal{C}' & \xrightarrow{\text{act}^*} & \mathcal{C}' \times_{S, p_1} \mathcal{G} \\ \text{act}^* \downarrow & & \text{act}^* \times \text{id}_{\mathcal{G}} \downarrow \\ \mathcal{C}' \times_{S, p_1} \mathcal{G} & \xrightarrow{\text{mult}^*} & \mathcal{C}' \times_{S, p_1} \left(\mathcal{G} \times_{p_2, S, p_1} \mathcal{G} \right) \end{array}$$

such that the natural compatibility conditions hold.

Lemma 14.

(a) Let $0 \rightarrow \mathcal{F}_1 \rightarrow \mathcal{F}_2 \rightarrow \mathcal{F}_3 \rightarrow 0$ be a short exact sequence of quasi-coherent sheaves on $\mathcal{G}$ with $\mathcal{F}_3$ being $\mathcal{O}_S$ -flat with respect to p_2 . Then for $X \in \mathcal{C}'$, the sequence

$$0 \rightarrow \mathcal{F}_1 * \text{act}^*(X) \rightarrow \mathcal{F}_2 * \text{act}^*(X) \rightarrow \mathcal{F}_3 * \text{act}^*(X) \rightarrow 0$$

is also short exact.

(b) If $X \in \mathcal{C}'$ is $\mathcal{O}_S$ -flat, then $\text{act}^*(X)$ is $\mathcal{O}_{\mathcal{G}}$ -flat.

Proof. Let S' be a scheme with a map $\phi : S' \rightarrow \mathcal{G}$; let $\psi_i = p_i \circ \phi$, $i = 1, 2$. We claim that there exists a natural $\mathcal{O}_{S'}$ -linear equivalence

$$\text{act}_{\phi}^* : \mathcal{C}' \times_{S, \psi_2} S' \rightarrow \mathcal{C}' \times_{S, \psi_1} S',$$

defined by

$$X \mapsto (\phi \times \text{id}_{S'})^* \circ (\text{act}^* \times \text{id}_{S'})(X),$$

where $\phi \times \text{id}_{S'} : S' \rightarrow \mathcal{G} \times_{p_2, S, \psi_2} S'$. Its quasi-inverse is defined using the map $\gamma \circ \phi : S' \rightarrow \mathcal{G}$, where γ is the inversion on $\mathcal{G}$.

We apply this to $S' = \mathcal{G}$ and $\phi = \gamma$. We obtain an equivalence

$$\text{act}_{\gamma}^* : \mathcal{C}' \times_{S, p_1} \mathcal{G} \rightarrow \mathcal{C}' \times_{S, p_2} \mathcal{G},$$

such that for $X \in \mathcal{C}'$,

$$\text{act}_{\gamma}^*(\text{act}^*(X)) \simeq p_2^*(X).$$

This readily implies both points of the lemma. □

We say that an object $X \in \mathcal{C}'$ is $\mathcal{G}$ -equivariant, if we are given an isomorphism

$$p_1^*(X) \simeq \text{act}^*(X) \in \mathcal{C}' \times_{S, p_1} \mathcal{G},$$

compatible with the unit and associativity constraints. Let us denote by $\mathcal{C}'^{\mathcal{G}}$ the category of $\mathcal{G}$ -equivariant objects in $\mathcal{C}'$.

From the definitions we obtain:

Lemma 15.

(a) For any $X \in \mathcal{C}'$, the object $(p_1)_*(\text{act}^*(X))$ is naturally $\mathcal{G}$ -equivariant.

(b) The functor $X \mapsto (p_1)_*(\text{act}^*(X))$ is the right adjoint to the forgetful functor $\mathcal{C}'^{\mathcal{G}} \rightarrow \mathcal{C}'$.

In addition, we have:

Lemma 16. Assume that $\mathcal{G}$ is flat over $S \times S$. Then every $\mathcal{G}$ -equivariant object of $\mathcal{C}'$ is $\mathcal{O}_S$ -flat.

Proof. This follows from the fact that for $\mathcal{F} \in \mathrm{QCoh}_S$ and $X \in \mathcal{C}'$,

$$\mathrm{act}^*(\mathcal{F} * X) \simeq p_2^*(\mathcal{F}) * \mathrm{act}^*(X).$$

□

17. Stacks: approach II. Let $\mathrm{Vect}_{\mathcal{Y}}$ denote the tensor category of locally free sheaves of finite rank on $\mathcal{Y}$.

Assume now that we are given a category $\mathcal{C}_{\mathcal{Y}}$ endowed with an action of the tensor category $\mathrm{Vect}_{\mathcal{Y}}$:

$$* : \mathrm{Vect}_{\mathcal{Y}} \times \mathcal{C}_{\mathcal{Y}} \rightarrow \mathcal{C}_{\mathcal{Y}},$$

which is exact. I.e., for a fixed $\mathcal{P} \in \mathrm{Vect}_{\mathcal{Y}}$ the functor $X \mapsto \mathcal{P} * X$ is exact, and whenever $0 \rightarrow \mathcal{P}_1 \rightarrow \mathcal{P}_2 \rightarrow \mathcal{P}_3 \rightarrow 0$ is a short exact sequence of objects of $\mathrm{Vect}_{\mathcal{Y}}$, the corresponding sequence

$$0 \rightarrow \mathcal{P}_1 * X \rightarrow \mathcal{P}_2 * X \rightarrow \mathcal{P}_3 * X \rightarrow 0$$

is also exact. We shall call such a data "a category over $\mathcal{Y}$ ".

We will now make an additional assumption on the stack $\mathcal{Y}$:

- The stack $\mathcal{Y}$ is locally Noetherian and every quasi-coherent sheaf on it is an inductive limit of coherent ones.
- Every coherent sheaf on $\mathcal{Y}$ can be covered by an object of $\mathrm{Vect}_{\mathcal{Y}}$.

As in the affine case, this implies that every flat quasi-coherent sheaf on $\mathcal{Y}$ can be represented as an inductive limit of objects of $\mathrm{Vect}_{\mathcal{Y}}$.

Theorem 18. *Under the above assumption on $\mathcal{Y}$, a data of a category over $\mathcal{Y}$ is equivalent to that of a sheaf of categories over $\mathcal{Y}$.*

The rest of this subsection and the next one are devoted to the proof of this theorem. One direction has been explained above: given a sheaf of categories $\mathcal{C}^{sh}$ over $\mathcal{Y}$, we reconstruct $\mathcal{C}_{\mathcal{Y}}$ as $\Gamma(\mathcal{Y}, \mathcal{C}^{sh})$. To carry out the construction in the opposite direction we will use the above additional assumption on $\mathcal{Y}$.

We claim that the above data extends to an action of the monoidal category $\mathrm{QCoh}_{\mathcal{Y}}$ on $\mathcal{C}_{\mathcal{Y}}$, satisfying the conditions (i),(ii),(iii) of Sect. 9 and assertions (a), (b) and (c) of Lemma 4.

First we define an action of the monoidal category $\mathrm{Coh}_{\mathcal{Y}}$ on $\mathcal{C}_{\mathcal{Y}}$: By assumption, every $\mathcal{F} \in \mathrm{Coh}_{\mathcal{Y}}$ can be represented as $\mathrm{coker}(\mathcal{P} \rightarrow \mathcal{Q})$ with $\mathcal{P}, \mathcal{Q} \in \mathrm{Vect}_{\mathcal{Y}}$. We set

$$\mathcal{F} * X := \mathrm{coker}(\mathcal{P} * X \rightarrow \mathcal{Q} * X).$$

To show that this is well-defined, we must consider a commutative diagram of objects of $\mathrm{Vect}_{\mathcal{Y}}$

$$\begin{array}{ccccccc} & 0 & & 0 & & 0 & \\ & \uparrow & & \uparrow & & \uparrow & \\ \mathcal{P} & \xrightarrow{\phi} & \mathcal{Q} & \longrightarrow & \mathcal{F} & \longrightarrow & 0 \\ & \uparrow & & \uparrow & & \mathrm{id} \uparrow & \\ \mathcal{P}' & \xrightarrow{\phi'} & \mathcal{Q}' & \longrightarrow & \mathcal{F} & \longrightarrow & 0 \\ & \uparrow & & \uparrow & & \uparrow & \\ \mathcal{P}'' & \xrightarrow{\phi''} & \mathcal{Q}'' & \longrightarrow & 0 & & \end{array}$$

with exact rows and columns, and show that the map

$$\mathrm{coker}(\phi * \mathrm{id}_X) \rightarrow \mathrm{coker}(\phi' * \mathrm{id}_X)$$

is an isomorphism. But this follows from the assumption.

It is clear that the resulting functor is right-exact and satisfies properties (ii) and (iii) of Sect. 9.

Next, we have to extend the above action of $\mathrm{Coh}_{\mathcal{Y}}$ on $\mathcal{C}_{\mathcal{Y}}$ to that of $\mathrm{QCoh}_{\mathcal{Y}}$ by setting for $\mathcal{F} \simeq \varinjlim \mathcal{F}_i$ with $\mathcal{F}_i \in \mathrm{Coh}_{\mathcal{Y}}$, $\mathcal{F} \in \mathrm{QCoh}_{\mathcal{Y}}$,

$$\mathcal{F} * X := \varinjlim \mathcal{F}_i * X.$$

The resulting action satisfies properties (i), (ii) and (iii) of Sect. 9. By assumption, the functor of tensor product with an object of $\mathrm{Vect}_{\mathcal{Y}}$ is exact. This implies properties (a) and (b) Lemma 4, by repeating the proof of *loc.cit.* Property (c) stated in Lemma 4 is evident.

19. We shall now show how the data of an action of $\mathrm{QCoh}_{\mathcal{Y}}$ on $\mathcal{C}_{\mathcal{Y}}$ with the above properties reconstructs the categories $\mathcal{C}_S$ for $S \in \mathrm{Sch}_{\mathcal{Y}}^{aff}$.

Let $S = \mathrm{Spec}(A)$, let us denote by $\mathcal{O}_S$ the direct image of the structure sheaf of S onto $\mathcal{Y}$, regarded as an algebra in $\mathrm{QCoh}_{\mathcal{Y}}$. We introduce $\mathcal{C}_S$ as the category, consisting of objects X of $\mathcal{C}_{\mathcal{Y}}$, endowed with an associative action $\mathcal{O}_S * X \rightarrow X$, and the morphisms being $\mathcal{C}_{\mathcal{Y}}$ -morphisms compatible with the action.

We have a map of algebras $A \otimes \mathcal{O}_{\mathcal{Y}} \rightarrow \mathcal{O}_S$ in $\mathrm{QCoh}_{\mathcal{Y}}$; this makes $\mathcal{O}_S$ into an A -linear category. We also have a functor $\mathcal{C}_{\mathcal{Y}} \rightarrow \mathcal{C}_S$ given by $X \mapsto X_S := \mathcal{O}_S * X$.

Let $f : S_2 = \mathrm{Spec}(A_2) \rightarrow \mathrm{Spec}(A_1) = S_1$ be a morphism in $\mathrm{Sch}_{\mathcal{Y}}^{aff}$. We define a functor $f^* : \mathcal{C}_{S_1} \rightarrow \mathcal{C}_{S_2}$ by

$$X \mapsto \mathcal{O}_{S_2} \underset{\mathcal{O}_{S_1}}{*} X,$$

where for an algebra $\mathcal{A}$ in $\mathrm{QCoh}_{\mathcal{Y}}$, a sheaf $\mathcal{F}$ of $\mathcal{A}$ -modules and an object $X \in \mathcal{C}_{\mathcal{Y}}$ acted on by $\mathcal{A}$, we set

$$\mathcal{F} \underset{\mathcal{A}}{*} X := \mathrm{coker}((\mathcal{F} \underset{\mathcal{O}_{\mathcal{Y}}}{*} X \rightrightarrows \mathcal{F} * X).$$

We claim that the induced functor $\mathcal{C}_{S_1} \times_{S_1} S_2 \rightarrow \mathcal{C}_{S_2}$ is an equivalence. This follows from the fact that $\mathcal{O}_{S_2} \simeq A_2 \underset{A_1}{\otimes} \mathcal{O}_{S_1}$.

Note in addition that for $X \in \mathcal{C}_{\mathcal{Y}}$,

$$\mathcal{O}_{S_2} * X \simeq A_2 \underset{A_1}{\otimes} (\mathcal{O}_{S_1} * X).$$

This implies that for $X \in \mathcal{C}_{\mathcal{Y}}$, we have a natural isomorphism $f^*(X_{S_1}) \simeq X_{S_2}$.

Thus, we have constructed a sheaf of categories over $\mathcal{Y}$, and it remains to show that the initial category $\mathcal{C}_{\mathcal{Y}}$ can be reconstructed by the descent procedure. The usual proof for coherent sheaves works, once we establish the following:

Lemma 20. *If $\mathcal{A} \in \mathrm{QCoh}_{\mathcal{Y}}$ be an algebra, faithfully flat over $\mathcal{O}_{\mathcal{Y}}$. Then the functor $X \mapsto \mathcal{A} * X : \mathcal{C}_{\mathcal{Y}} \rightarrow \mathcal{C}_{\mathcal{Y}}$ is exact and faithful.*

Proof. The exactness part follows by property (a) of Lemma 4. The faithfulness part follows as in Proposition 5 using property (b) of Lemma 4. $\square$

21. Example: de-equivariantization. Let $\mathcal{Y}$ be the stack pt/G , where G is an affine algebraic group. Given a category $\mathcal{C}$, a structure of category over pt/G on it is by definition the same as an action of the tensor category $\text{Rep}(G)$ of finite-dimensional representations of G on it:

$$V \in \text{Rep}(G), X \in \mathcal{C} \mapsto V * X \in \mathcal{C},$$

which has the exactness property of Sect. 17.

By Theorem 18, such a data gives rise to a sheaf of categories $\mathcal{C}^{sh}$ over pt/G .

Let us show how to reconstruct the category $\mathcal{C}' := \mathcal{C}^{sh} \times_{\text{pt}/G} \text{pt}$. By definition, this is the category, whose objects are $X' \in \mathcal{C}$, endowed with an associative action $\mathcal{O}_G * X' \rightarrow X'$, and morphisms are $\mathcal{C}$ -morphisms, compatible with this action.

According to [AG], this data can be rewritten as follows. For every $V \in \text{Rep}(G)$ we must be given a map

$$\beta_V : V * X \rightarrow X \otimes \underline{V},$$

for every $V \in \text{Rep}(G)$ (here $\underline{V}$ denoted the vector space underlying a representation), which satisfy the compatibility conditions of [AG], Sect. 2.2. One easily shows that the maps β_V are necessarily isomorphisms. Morphisms in this category are $\mathcal{C}$ -morphisms, compatible with the data of β .

Thus, $\mathcal{C}'$ is the category of Hecke eigen-objects in $\mathcal{C}$ with respect to the action of $\text{Rep}(G)$. By construction, $\mathcal{C}'$ carries a canonical action of G . Explicitly, for $X' \in \mathcal{C}'$, the $\mathcal{O}_G$ -family $\text{act}^*(X')$ is isomorphic to $X' \otimes \mathcal{O}_G$ as an object of $\mathcal{C}$. The isomorphisms β are given by

$$V * (X' \otimes \mathcal{O}_G) \xrightarrow{\beta_V \otimes \text{id}_{\mathcal{O}_G}} X' \otimes \underline{V} \otimes \mathcal{O}_G \rightarrow X' \otimes \underline{V} \otimes \mathcal{O}_G,$$

where the second arrow is given by the co-action map $\underline{V} \rightarrow \underline{V} \otimes \mathcal{O}_G$.

According to Sect. 6, the category $\mathcal{C}$ is reconstructed from $\mathcal{C}'$ as the category $\mathcal{C}'^G$ of G -equivariant objects. We will refer to $\mathcal{C}'$ as the de-equivariantization of $\mathcal{C}$.

22. Another example Generalizing the previous example, let us take $\mathcal{Y} = S/G$, where $S = \text{Spec}(A)$ is an affine scheme, and G an affine algebraic group acting on it. Let $\mathcal{C}$ be an abelian category. A structure on $\mathcal{C}$ of category over S/G is by definition expressed as follows:

An action of the tensor category $\text{Rep}(G)$ on $\mathcal{C}$: $V, X \mapsto V * X$ as above, and a functorial map $\alpha_X : A * X \rightarrow X$, where A is regarded as an algebra in $\text{Rep}(G)$, such that for $V \in \text{Rep}(G)$ the diagram

$$\begin{array}{ccc} V * (A * X) & \xrightarrow{\sim} & A * (V * X) \\ V * \alpha_X \downarrow & & \alpha_{V * X} \downarrow \\ V * X & \xrightarrow{\text{id}} & V * X \end{array}$$

commutes.

REFERENCES

- [AG] S. Arkhipov and D. Gaiitsgory, *Another realization of the category of modules over the small quantum group*, math.QA/0010270, Adv. Math. **173** (2003), 114–143.

DEPARTMENT OF MATHEMATICS, HARVARD UNIVERSITY, 1 OXFORD STREET, CAMBRIDGE MA, 02138
E-mail address: gaiitsgde@math.harvard.edu